

(12) UK Patent Application (19) GB (11) 2 395 270 (13) A

(43) Date of A Publication 19.05.2004

(21) Application No: 0226622.9

(22) Date of Filing: 14.11.2002

(71) Applicant(s):
The University Of Nottingham
(Incorporated in the United Kingdom)
University Park, NOTTINGHAM, NG7 2RD,
United Kingdom

(72) Inventor(s):
Catherine Rosamund Louise Graves
John Forsyth Russell Robertson

(74) Agent and/or Address for Service:
Boult Wade Tennant
Verulam Gardens, 70 Gray's Inn Road,
LONDON, WC1X 8BT, United Kingdom

(51) INT CL⁷:
G01N 33/574 , A61P 35/00

(52) UK CL (Edition W):
G1B BAE BAX B300 B402
U1S S1313

(56) Documents Cited:
WO 2001/011372 A1 **WO 2000/026668 A2**
WO 1999/058978 A2
Clin Cancer Res; Vol 7, pp 781s-787s (2001). Beatty et
al. "Biochemical characterisation of the soluble
form..."
Lung; Vol 170, pp 65-74 (1992). Lawniczak et al. "The
search for tumour-associated proteins in pleural
effusions..."
Clin Chim Acta; Vol 300, pp 43-55 (2000). Kuralay et al.
"Diagnostic usefulness of tumour marker levels..."
Intern Med; Vol 36, pp 724-727 (1997). Yamamoto et
al. "L-Myc overexpression and detection of
auto-antibodies..."

(continued on next page)

(54) Abstract Title: **METHOD OF DETECTION OF TUMOUR-ASSOCIATED AUTOANTIBODIES**

(57) An immunoassay for detecting cancer-associated anti-tumour autoantibodies comprising contacting a sample with a tumour marker protein. The tumour marker protein is prepared from the bodily fluid, excretion, or derived from a body cavity or space in which a tumour is or was associated, or one or more cancer patients. The method can be used in the diagnosis of cancer. In an alternative embodiment there is disclosed methods for isolating tumour markers from bodily fluids collected from a body cavity or space, and another embodiment relates to the preparation of tumour marker proteins from an excretion.

GB 2 395 270 A

(56) cont

Int J Oncology; Vol 13, pp 605-610 (1998). Monternarh et al. "p53 autoantibodies in the sera, cyst and ascitic fluids..."

Br J Cancer; Vol 80, pp 1987-1994 (1999). Schneider et al. "p53 protein, EGF receptor and anti-p53 antibodies..."

Clin Cancer Res; Vol 4, pp 3025-3030 (1998). Lai et al. "Presence of serum anti-p53 antibodies..."

(58) Field of Search:

INT CL⁷ **G01N**

Other: **WPI, EPODOC, JAPIO, MEDLINE, EMBASE, BIOSIS, SCISEARCH, CAPLUS**

FIG. 1 Post Ig disruption gel filtration chromatogram.





FIG. 2.


Silver stained gel, c-myc purification, post immunoaffinity.



- 1 – major c-myc peak, room temp purification
- 2 – major c-myc peak, 4°C purification
- 3 – minor c-myc peak, room temp purification
- 4 – minor c-myc peak, 4°C purification

FIG. 3.

Immunoprobed blot, c-myc purification, post immunoaffinity.



- 1 – major c-myc peak, room temp purification
- 2 – major c-myc peak, 4°C purification
- 3 – minor c-myc peak, room temp purification
- 4 – minor c-myc peak, 4°C purification
- 5 – molecular weight markers

FIG. 4.

Serum Samples vs MUC1 from Various Body Fluids.




FIG. 5.

Normal Serum Samples vs MUC1 from Various Body Fluids.




FIG. 6.

Serum samples vs normal and tumour associated CA125





FIG. 7.

Concentration of MUC1 in sera and fluids derived from individuals with cancer.




PE – pleural effusion.

AF – ascitic fluid.

FIG. 8.


Concentration of MUC1 in serum and fluids derived from an individual with cancer.



WDF – wound draining fluid

FIG. 9.

Reactivity of human anti-MUC1 antibodies purified against cancer associated MUC1 from seroma.



Pt-nMUC1 – urinary MUC1 from patient M, 2 years prior to cancer diagnosis.
Pt-MUC1 – MUC1 derived from the seroma of patient M, after diagnosis with cancer.
BSA-AG – bovine serum albumen conjugated to MUC1 protein core peptide.

FIG. 10.

Serum reactivity against MUC1 purified from Ascitic Fluid.






FIG. 11.

Serum reactivity against MUC1 purified from Pleural Effusions.



12/12
FIG. 12.
Titration of PE MUC1



Tumour Marker Proteins and Uses ThereofField of the invention

5 The invention relates to tumour marker proteins and their preparation from fluids from one or more cancer patients wherein said fluids are those which collect in a body cavity or space which is naturally occurring or which is the result of cancer or medical intervention for cancer. Exemplary fluids are
10 ascites, pleural effusion, seroma, hydrocoele and wound drainage fluid. The invention also relates to preparation of tumour marker proteins from excretions taken from patients with cancer.

15 The said tumour marker proteins are useful in cancer detection methods which involve detecting or quantitatively measuring autoantibodies to circulating tumour markers or markers expressed on or in tumour cells and in various research applications. The
20 invention is also directed to such uses.

Background to the invention

The development and progression of cancer in a patient is generally found to be associated with the
25 presence of markers in the bodily fluid of the patient, these "tumour markers" reflecting different aspects of the biology of the cancer (see Fateh-Maghadam, A. & Steilber, P. (1993) Sensible use of tumour markers. Published by Verlag GMBH, ISBN 3-
30 926725-07-9). Tumour markers are often found to be altered forms of wild-type proteins expressed by "normal" cells, in which case the alteration may be a change in primary amino acid sequence, a change in secondary, tertiary or quaternary structure or a
35 change in post-translational modification, for example, abnormal glycosylation. In addition, wild-type proteins which are up-regulated or over-expressed

in tumour cells, possibly as a result of gene amplification or abnormal transcriptional regulation, may also be tumour markers.

5 Established assays for tumour markers present in
bodily fluids tend to focus on the detection of tumour
markers which reflect tumour bulk and as such are of
value late in the disease process, for example in the
diagnosis of metastatic disease. The most widely used
10 of these markers include carcinoembryonic antigen
(CEA) and the glycoprotein termed CA 15.3, both of
which have been useful mainly as indicators of
systemic disease burden and of relapse following
therapy (Molina, R., Zanon, G., Filella, X. et al. Use
15 of serial carcinoembryonic antigen and CA 15.3 assays
in detecting relapses in breast cancer patients.
(1995) *Breast Cancer Res Treat* 36: 41-48). These
markers are of limited use earlier in the course of
the disease, for example in early detection or in the
20 screening of asymptomatic patients. Thus, in the
search for tumour markers present in bodily fluid that
are of use in assisting diagnosis earlier in the
disease process the present inventors have sought to
identify markers which do not depend on tumour bulk
25 *per se*.

Differences between a wild type protein expressed
by "normal" cells and a corresponding tumour marker
protein may, in some instances, lead to the tumour
marker protein being recognised by an individual's
30 immune system as "non-self" and thus eliciting an
immune response in that individual. This may be a
humoral (i.e B cell-mediated) immune response leading
to the production of autoantibodies immunologically
specific to the tumour marker protein. Autoantibodies
35 are naturally occurring antibodies directed to an
antigen which an individual's immune system recognises

as foreign even though that antigen actually
originated in the individual. They may be present in
the circulation as circulating free autoantibodies or
in the form of circulating immune complexes consisting
5 of autoantibodies bound to their target tumour marker
protein.

As an alternative to the direct measurement or
detection of tumour marker protein in bodily fluids,
10 assays may be developed to measure the immune response
of the individual to the presence of tumour marker
protein in terms of autoantibody production. Such
assays essentially constitute indirect detection of
the presence of tumour marker protein. Because of the
15 nature of the immune response, it is likely that
autoantibodies can be elicited by a very small amount
of circulating tumour marker protein and indirect
methods which rely on detecting the immune response to
tumour markers will consequently be more sensitive
20 than methods for the direct measurement of tumour
markers in bodily fluids. Assay methods based on the
detection of autoantibodies may therefore be of
particular value early in the disease process and
possibly also in relation to screening of asymptomatic
25 patients, for example in screening to identify
individuals "at risk" of developing disease amongst a
population of asymptomatic individuals. Furthermore,
they may be useful for earlier detection of recurrent
disease.

30 Tumour marker proteins observed to elicit serum
autoantibodies include a particular class of mutant
p53 protein, described in US Patent No. 5,652,115,
which can be defined by its ability to bind to the 70
35 kd heat shock protein (hsp70). p53 autoantibodies can
be detected in patients with a number of different
benign and malignant conditions (described in US

5,652,115) but are in each case present in only a subset of patients. For example, one study utilizing an ELISA assay for detection of autoantibodies directed against the p53 protein in the serum of breast cancer patients reported that p53 autoantibodies were produced by 26% of patients and 1.3% of control subjects (Mudenda, B., Green, J. A., Green, B. et al. The relationship between serum p53 autoantibodies and characteristics of human breast cancer, (1994) *Br J Cancer* **69**: 4445-4449). A second tumour marker protein known to elicit serum autoantibodies is the epithelial mucin MUC1 (Hinoda, Y. et al. (1993) *Immunol Lett.* 35: 163-168; Kotera, Y. et al. (1994) *Cancer Res.* 54: 2856-2860).

WO 99/58978 describes methods for use in the detection/diagnosis of cancer which are based on evaluating the immune response of an individual to two or more distinct tumour markers. These methods generally involve contacting a sample of bodily fluid taken from the individual with a panel of two or more distinct tumour marker antigens, each derived from a separate tumour marker protein, and detecting the formation of complexes of the tumour marker antigens bound to circulating autoantibodies immunologically specific for the tumour marker proteins. The presence of such circulating autoantibodies is taken as an indication of the presence of cancer.

Cancer detection methods based on detection of circulating autoantibodies are frequently immunoassays utilizing an "immunoassay reagent" reactive with the circulating autoantibodies. Typically, the "reagents" used in such assays comprise recombinant tumour marker proteins (expressed in bacterial, insect, yeast or mammalian cells) or chemically synthesised tumour

marker antigens, which may comprise substantially whole tumour marker proteins, or fragments thereof, such as short peptide antigens. Other potential sources of tumour-associated proteins for use as the
5 basis of immunoassay reagents for the detection of anti-tumour auto-antibodies include cultured tumour cells (and the spent media used for their growth), tumour tissue, and serum from individuals with neoplasia. The majority of these sources have
10 significant drawbacks, as discussed below.

With cultured tumour cells (and their spent media) the amount of expressed protein can vary depending on growth phase at the time of harvest,
15 leading to variations in quality and quantity. In addition, the desired protein is generally present at low concentration, therefore it is time-consuming to purify sufficient quantities of protein. Furthermore, the cell stock will be clonal, unlike cell stock in a
20 tumour which is likely to have become heterogeneous in nature during the growth of the neoplasm, therefore producing variations in protein (especially in the degree of glycosylation).

25 Recombinant proteins expressed in bacterial cells are not glycosylated, and thus significantly different from naturally glycosylated proteins. In addition, refolding of recombinantly expressed proteins may not be appropriate, thus giving an incorrect conformation
30 for auto-antibody recognition.

Tumour tissue is usually only available in small quantities and the purification of proteins therefrom is laborious and time consuming.

35 Serum samples are usually available only in small quantities, therefore it is difficult to purify

sufficient quantities of protein.

5 The present inventors have now determined that significant advantages can be gained by the use of tumour marker antigens purified from bodily fluids derived from a body cavity or space in which a tumour is present or with which it is or was associated, such as ascites fluid, pleural effusion, seroma, hydrocoele or wound drainage fluid, or from excretions, as the
10 "reagent" in auto-antibody immunoassays. In particular, the inventors have observed that use of reagents comprising tumour marker antigens purified from bodily fluids derived from the above defined body cavities or spaces results in increased
15 sensitivity (as compared to the use of reagents derived from a "normal" body fluid) and produces a more "clinically relevant" result. There are also significant practical advantages to be gained from the use of such fluids as a source of assay reagent.

20

Summary of the invention

In a first aspect the invention relates to a method of detecting cancer-associated anti-tumour autoantibodies, which method is an immunoassay
25 comprising contacting a sample to be tested for the presence of such autoantibodies with an immunoassay reagent and detecting the presence of complexes formed by specific binding of the immunoassay reagent to any cancer-associated anti-tumour autoantibodies present
30 in the sample, wherein the immunoassay reagent comprises tumour marker protein prepared from bodily fluid derived from a body cavity or space within which a tumour is or was present or with which a tumour is or was associated, from one or more cancer patients
35 and/or tumour marker protein prepared from an excretion from one or more cancer patients, wherein said tumour marker protein exhibits selective

reactivity with cancer-associated anti-tumour autoantibodies.

5 In a second aspect the invention relates to use of tumour marker protein prepared from bodily fluid derived from a body cavity or space within which a tumour is or was present or with which a tumour is or was associated, of one or more cancer patients and/or tumour marker protein derived from an excretion of one or more cancer patients in the manufacture of an immunoassay reagent exhibiting selective reactivity with cancer-associated anti-tumour autoantibodies.

15 In a third aspect, the invention relates to a method of preparing a tumour marker protein which method comprises isolating said tumour marker protein from bodily fluid wherein said fluid is:

- 20 (i) collected from a body cavity or space in which a tumour is or was present or with which a tumour is or was associated, and
- (ii) said fluid represents the pooled fluid samples from two or more cancer patients.

25 In a fourth aspect, the invention relates to tumour marker protein preparations prepared as described above which are substantially immunoglobulin free and to kits and reagents comprising said preparations.

30 Detailed description of the invention

In the first aspect, the invention relates to a method of detecting "cancer-associated" anti-tumour autoantibodies.

35 The term "cancer-associated" anti-tumour autoantibodies refers to autoantibodies which are characteristic of the cancer disease state, and which

are directed against epitopes present on forms of tumour marker proteins which are preferentially expressed in the cancer disease state.

5 The method of the invention comprises an immunoassay to detect and/or quantitatively measure autoantibodies immunologically specific for one or more tumour marker proteins, and is characterised in that the "immunoassay reagent" used in the immunoassay
10 comprises tumour marker protein prepared from bodily fluid derived from a body cavity or space in which a tumour is or was present or with which a tumour is or was associated, from one or more cancer patients and/or tumour marker protein prepared from an
15 excretion of one or more cancer patients. Generally, excretion will be through an organ in which cancer is present wherein the excretion is in contact with said cancer, or the excretion will include one or more components which have been in contact with cancer
20 elsewhere in the body. A particular example is bile which may be in contact with cancer in the gall bladder but will appear in the faeces.

 The immunoassay reagent exhibits "selective reactivity" with cancer-associated anti-tumour
25 autoantibodies. As used herein "selective reactivity" means a tumour marker protein has a greater affinity for autoantibodies to the tumour-associated antigen than it does for any antibody or autoantibody made to
30 the same antigen which exists in the normal i.e. non-tumour possessing state.

 The term "body cavity or space" includes any body cavity or space, whether it be a natural cavity or a
35 space or cavity arising as a result of diseases or medical intervention including collapsed or former cavities. The fluid is derived from such a cavity or

space in which a tumour is or was present or with which a tumour is or was associated. Preferably the "bodily fluid derived from a body cavity" will be a tumour-induced body fluid, meaning a body fluid which is produced during the disease process, for example in response to or as a consequence of the presence of tumour cells. Exemplary body fluids are ascites, pleural effusion, seroma, hydrocoele and wound drainage fluid.

For the avoidance of doubt "bodily fluids derived from a body cavity or space" do not include fluids derived from the systemic circulation, such as whole blood or serum.

The term "excretion" includes, *inter alia*, urine, faeces, and seminal fluid.

The general features of immunoassays, for example ELISA, radioimmunoassays and the like, are well known to those skilled in the art (see Immunoassay, E. Diamandis and T. Christopoulos, Academic Press, Inc., San Diego, CA, 1996). Immunoassays for the detection of antibodies having a particular immunological specificity (e.g. autoantibodies having immunological reactivity with a given tumour marker protein) generally require the use of a reagent that exhibits specific immunological reactivity with the antibody under test. Depending on the format of the assay this reagent may be immobilised on a solid support. A sample to be tested for the presence of the antibody is brought into contact with the reagent and if antibodies of the required immunological reactivity are present in the sample they will immunologically react with the reagent to form autoantibody-reagent complexes which may then be detected or quantitatively measured.

Suitable samples of tumour marker protein for use as the basis of the "immunoassay reagent" may be isolated from bodily fluids derived from a body cavity or space from one or more cancer patients and/or from excretions from one or more cancer patients using standard protein purification techniques, such as are generally known in the art. For example, tumour marker proteins may be isolated by affinity chromatography using a suitable antibody (or antibody fragment) immunologically specific for the tumour marker protein.

The starting material of bodily fluids derived from a body cavity and/or excretions is/are taken from one or more cancer patients. In this context the term "cancer patient" includes an individual previously diagnosed as having cancer. The fluid/excretion may be taken from a single patient or samples from two or more patients may be pooled together. Samples may be pooled from two or more patients having the same or different stages of the same or different types of cancers. Samples may also be pooled from different types of bodily fluids or excretions from a single or multiple patients. Advantageously, an immunoassay reagent prepared from fluid and/or excretion taken from cancer patient(s) with a particular type of cancer may be used to assist in the diagnosis of the same types of cancers in other individuals.

In one embodiment the "cancer patient" from which the fluid/excretion is taken may be the same patient which it is later intended to test using the assay reagent. For example, a stock of reagent prepared from a patient diagnosed with cancer may be used at a later date to assess the immune status of the same patient, for example to monitor disease progression and/or to assess the effectiveness of a course of

anti-cancer treatment in that patient.

5 The "immunoassay reagent" or "tumour marker
preparation" may comprise substantially whole tumour
marker protein, for example tumour marker protein
substantially in the form in which it is isolated from
the fluid/excretion, or it may comprise a fragment of
the tumour marker protein. To be effective as an
10 immunoassay reagent any such "fragment" must retain
immunological reactivity with the (auto)antibodies for
which it is desired to test using the reagent.
Suitable fragments might, for example, be prepared by
chemical or enzymatic cleavage of the isolated tumour
marker protein.

15 Depending on the precise nature of the
immunoassay in which it will be used, the "reagent" or
"tumour marker protein preparation" may comprise a
tumour marker protein, or fragment thereof, linked to
20 one or more further molecules which impart some
desirable characteristic not naturally present in the
tumour marker protein. For example, the tumour marker
protein may be conjugated to a revealing label, such
as a fluorescent label, coloured label, luminescent
25 label, radiolabel or heavy metal such as colloidal
gold.

30 The tumour marker protein as prepared by the
method described herein can also be immobilized for
use on a solid support such as a bead or surface of a
well of a multiwell plate. The immobilization may be
by absorption or by co-valent attachment. The tumour
marker protein is preferably substantially
35 immunoglobulin free by virtue of the fact that
following isolation, for example, by affinity
chromatography, the protein preparation is treated to
specifically remove contaminating immunoglobulins.

The use of an immunoassay reagent comprising a tumour marker protein (or fragment thereof) isolated from body cavity fluids and/or excretions taken from one or more cancer patients provides significant advantages over the use of other reagents, such as recombinantly expressed or chemically synthesised polypeptides, in the clinical detection of cancer (including diagnosis, monitoring of disease recurrence or disease progression, etc).

10

It might be expected that the precise characteristics of tumour marker proteins isolated from cancer patients could vary depending upon the source material (e.g. tissue or fluid) from which the tumour marker protein is isolated. For example, the characteristics of proteins isolated from urine may be different to those isolated from whole blood or serum, which may be different again to those isolated from ascites or pleural effusion. This may in turn affect the utility of the tumour marker protein as an assay reagent.

20

In fact, the inventors have surprisingly observed that reagents prepared from tumour marker proteins isolated from body cavity-derived fluids or excretions from cancer patients, particularly ascites fluid, pleural effusion, seroma or wound drainage fluid are generally more specific for cancer-associated autoantibodies than reagents based on the equivalent proteins isolated from "normal" individuals. This increased specificity for cancer-associated autoantibodies means that immunoassays based on the use of reagents prepared from body cavity-derived fluids or excretions from cancer patients produce results that are more "clinically relevant" in the detection of an immune response to cancer.

25

30

35

The inventors postulate that the improved specificity observed with the use reagents prepared from fluids derived from body of cavities of cancer patients, such as ascites, pleural effusion, seroma or wound drainage fluid, is due to the origin of such fluids within the body cavities or spaces of cancer patients. It is postulated that fluids originating in body cavities or spaces due to the presence of a tumour in contact with the major organs, may pick up more "cancer-associated" forms of the tumour marker protein, which are actually relevant to the cancer disease state. Since it is generally differences between "tumour" marker proteins and their "normal" counterparts which trigger the development of an immune response (i.e. autoantibody production), the inventors hypothesise that reagents based on the use of tumour markers isolated from cancer patients will be more specific for cancer autoantibodies than the equivalent "normal" proteins. This is indeed the case with tumour marker antigens isolated from ascites, pleural effusion or seroma, as shown in the accompanying Examples.

There are further practical advantages associated with the use of ascites fluid, pleural effusion, seroma, hydrocoele or wound drainage fluid, as a source of tumour marker proteins. These fluids may be readily removed from patients in relatively large volumes as part of the therapeutic strategy. This material, which would otherwise be discarded, is a valuable source of useful assay reagent.

Furthermore, it has also been observed by the inventors that additional significant advantages can be secured by pooling body cavity fluid samples from two or more patients. Apart from increasing protein yield, the product secures at least as good a

detection rate as marker protein from an individual sample while, at the same time, being more consistent in its characteristics from batch to batch. Thus, adequate affinity of the antigen can be relied upon every time.

In particular embodiments the methods of the invention may comprise immunoassays to (simultaneously) detect two or more types of autoantibodies, each having specificity for different tumour marker proteins or for different epitopes on the same tumour marker proteins. These methods will typically involve use of a panel of two or more assay reagents, each reagent comprising a different tumour marker protein. These methods, which may be hereinafter referred to as "panel assays", utilise a panel of two or more reagents to monitor the overall immune response of an individual to a tumour or other carcinogenic/neoplastic change. These methods thus detect a "profile" of the immune response in a given individual, indicating which tumour markers elicit an immune response resulting in autoantibody production. The use of a panel of two or more reagents to monitor production of autoantibodies against two or more different tumour markers is generally more sensitive than the detection of autoantibodies to single markers and gives a much lower frequency of false negative results.

The methods of the invention are preferred for the detection of circulating free autoantibodies, but may be adapted for detection of autoantibodies present in immune complexes, as would be appreciated by the skilled reader, for example by the competitive use of labelled tumour marker.

In preferred applications the method of the

invention will be used to detect the presence of cancer-associated anti-tumour autoantibodies in human subjects or patients, and will most preferably take the form of an *in vitro* immunoassay, performed on
5 samples of bodily fluid taken from the subject/patient. Such *in vitro* immunoassays are non-invasive and can be repeated as often as is thought necessary to build up a profile of autoantibody production in a patient, either prior to the onset of
10 disease, as in the screening of "at risk" individuals, or throughout the course of disease (further discussed below in relation to preferred applications of the method). As used herein the term "bodily fluid", when referring to the material to be tested for the
15 presence of autoantibodies by immunoassay, includes *inter alia* plasma, serum, whole blood, urine, sweat, lymph, faeces, cerebrospinal fluid, ascites, pleural effusion, seminal fluid, sputum or nipple aspirate. The type of bodily fluid used may vary depending upon
20 the type of cancer involved and the clinical situation in which the assay is used. In general, it is preferred to perform the assays on samples of serum or plasma.

25 As aforesaid, the "immunoassay" used to detect/quantitate cancer-associated autoantibodies may be carried out according to standard techniques known in the art. In a most preferred embodiment the immunoassay may be an ELISA. ELISAs are generally
30 well known in the art. In a typical "sandwich" ELISA a reagent having specificity for the autoantibodies under test is immobilised on a solid surface (e.g. the wells of a standard microtiter assay plate, or the surface of a microbead) and a sample of body fluid to
35 be tested for the presence of autoantibodies is brought into contact with the immobilised reagent. Any autoantibodies of the desired specificity present

in the sample will bind to the immobilised reagent. The bound autoantibody/reagent complexes may then be detected using any suitable method. In one preferred embodiment a labelled secondary anti-human
5 immunoglobulin antibody, which specifically recognises an epitope common to one or more classes of human immunoglobulins, is used to detect the autoantibody/reagent complexes. Typically the secondary antibody will be anti-IgG or anti-IgM. The
10 secondary antibody is usually labelled with a detectable marker, typically an enzyme marker such as, for example, peroxidase or alkaline phosphatase, allowing quantitative detection by the addition of a substrate for the enzyme which generates a detectable
15 product, for example a coloured, chemiluminescent or fluorescent product. Other types of detectable labels known in the art may be used with equivalent effect.

ELISA's may be performed in a qualitative format,
20 in which the objective is merely to determine the presence or absence of autoantibodies in the sample, or in a quantitative format, which provides a measurement of the quantity of autoantibodies present in the sample. For quantitative assays, a standard
25 curve may be generated by measuring the signal obtained (using the same detection reaction as will be used for the assay) from a series of standard samples containing known concentrations of antibodies having similar specificity as the autoantibodies under test.
30 The quantity of autoantibodies present in the sample under test may then be interpolated from the standard curve.

Panel assays may be performed in a multi-well
35 format in which each one of the two or more assay reagents is placed in a separate well of a multi-well assay plate or, alternatively, in a single-pot format

in which the two or more assay reagents are placed in a single container.

5 The method of the invention may be adapted for
use in the detection of autoantibodies to essentially
any tumour marker protein for which a suitable "assay
reagent" may be prepared from bodily fluid derived
from a body cavity and/or from an excretion from a
cancer patient. In particular, the method may be
10 adapted to detect/measure autoantibodies to the
epidermal growth factor receptor-related protein c-
erbB2 (Dsouza, B. et al. (1993) *Oncogene*. 8: 1797-
1806), the glycoprotein MUC1 (Batra, S. K. et al.
(1992) *Int. J. Pancreatology*. 12: 271-283) and the
15 signal transduction/cell cycle regulatory proteins Myc
(Blackwood, E. M. et al. (1994) *Molecular Biology of
the Cell* 5: 597-609), p53 (Matlashewski, G. et al.
(1984) *EMBO J.* 3: 3257-3262; Wolf, D. et al. (1985)
Mol. Cell. Biol. 5: 1887-1893) and ras (or Ras)
20 (Capella, G. et al. (1991) *Environ Health
Perspectives*. 93: 125-131), and also BRCA1 (Scully, R.
et al. (1997) *PNAS* 94: 5605-10), BRCA2 (Sharan, S. K.
et al. (1997) *Nature*. 386: 804-810), APC (Su, L. K. et
al. (1993) *Cancer Res.* 53: 2728-2731; Munemitsu, S.
25 et al. (1995) *PNAS* 92: 3046-50), CA125 (Nouwen, E. J.
et al. (1990) *Differentiation*. 45: 192-8), PSA
(Rosenberg, R. S. et al. (1998) *Biochem Biophys Res
Commun.* 248: 935-939), carcinoembryonic antigen CEA
(Duffy, M.J. (2001) *Clin Chem*, Apr 47(4):624-30), and
30 CA19.9 (Haga, Y. et al (1989) *Clin Biochem* (1989) Oct
22(5): 363-8). However, the invention is not intended
to be limited to the detection of autoantibodies to
these particular tumour markers.

35 The assay method of the invention may be employed
in a variety of different clinical situations. In

particular, the method may be used in the detection or diagnosis of cancer, in monitoring the progress of cancer or other neoplastic disease in a patient, in detecting early neoplastic or early carcinogenic
5 change in an asymptomatic human subject, in screening a population of asymptomatic human subjects in order to identify those subjects who are at increased risk of developing cancer, in monitoring the response of a cancer patient to anti-cancer treatment, in the
10 detection of recurrent disease in a patient previously diagnosed as having cancer who has undergone anti-cancer treatment to reduce the amount of cancer present, or in the selection of an anti-cancer vaccine for use in a particular patient.

15
The inventors have generally observed that levels of cancer-associated autoantibodies show a positive correlation with disease state (see also WO 99/58979, the contents of which are incorporated herein by
20 reference). Hence, when the method of the invention is used in clinical applications increased levels of anti-tumour marker autoantibodies, as compared to suitable controls, are generally taken as an indication of the cancer disease state.

25
For example, when the immunoassays are used in the diagnosis of cancer, the presence of an elevated level of autoantibodies, as compared to "normal" control individuals, is taken as an indication that
30 the individual has cancer. The "normal" control individuals will preferably be age-matched controls not having any diagnosis of cancer based on clinical, imaging and/or biochemical criteria.

35
When the immunoassays are used in monitoring the progress of cancer or other neoplastic disease in a patient, the presence of an elevated level of

autoantibodies, as compared to a "normal control", is taken as an indication of the presence of cancer in the patient. The "normal control" may be levels of autoantibodies present in control individuals, preferably age-matched, not having any diagnosis of cancer based on clinical, imaging and/or biochemical criteria. Alternatively, the "normal control" may be a "base-line" level established for the particular patient under test. The "base-line" level may be, for example, the level of autoantibodies present when either a first diagnosis of cancer or a diagnosis of recurrent cancer was made. Any increase above the base-line level would be taken as an indication that the amount of cancer present in the patient has increased, whereas any decrease below the base-line would be taken as an indication that the amount of cancer present in the patient has decreased. The "base-line" value may also be, for example, the level before a new treatment is commenced. A change in the level of autoantibodies would be taken as an indication of the effectiveness of the therapy. The direction of the "change" (i.e. increase vs decrease) indicating a positive response to treatment will be dependent upon the precise nature of the treatment. For any given treatment the direction of the "change" in autoantibody levels indicating a positive result may be readily determined, for example by monitoring autoantibody levels in comparison to other clinical or biochemical indicators of response to the treatment.

When the immunoassays are used in screening a population of asymptomatic human subjects to identify those subjects who are at increased risk of developing cancer, individuals having an elevated level of autoantibodies, as compared to "normal" control individuals, are identified as being "at risk" of developing cancer. The "normal" control individuals

will preferably be age-matched controls not identified as having any predisposition to developing cancer or any significant elevated risk of developing cancer. An exception to this may be where age itself is a
5 major risk factor.

When the immunoassays are used in monitoring the response of a cancer patient to anti-cancer treatment, the presence of a decreased level of autoantibodies
10 after treatment is taken as an indication that the patient has responded positively to the treatment. A base-line level of autoantibodies taken before treatment is commenced may be used for comparison purposes in order to determine whether treatment
15 results in a "decrease" in autoantibody levels.

When the immunoassays are used in detection of recurrent disease, the presence of an increased level of autoantibodies in the patient, as compared to a
20 "normal control", is taken as an indication that disease has recurred. The "normal control" may be levels of autoantibodies present in control individuals, preferably age-matched not having any diagnosis of cancer based on clinical, imaging and/or
25 biochemical criteria. Alternatively, the "normal control" may be a "base-line" level established for the particular patient under test. The "base-line" level may be, for example, the level of autoantibodies present during a period of remission from disease
30 based on clinical, imaging and/or biochemical criteria.

The assay method of the invention may be applied in the detection of many different types of cancer, of
35 which examples are breast, bladder, colorectal, prostate and ovarian cancers. The assays may complement existing methods of screening and

surveillance. For example, in the case of primary breast cancer immunoassays for autoantibodies could be used to alert clinicians to biopsy small lesions on mammograms which radiographically do not appear
5 suspicious or to carry out breast imaging or to repeat imaging earlier than planned. In the clinic, the assay methods of the invention are expected to be more objective and reproducible compared to current imaging techniques (i.e. mammography and ultrasound), the
10 success of which can be operator-dependent.

"Panel assays" may be tailored having regard to the particular clinical application. A panel of reagents for detection of autoantibodies to at least
15 p53 and c-erbB2 is particularly useful for many types of cancer and can optionally be supplemented with other markers having a known association with the particular cancer, or a stage of the particular cancer, to be detected. For example for breast cancer
20 the panel might include MUC 1 and /or c-myc and/or BRCA1 and/or BRCA2 and/or PSA whereas bladder cancer the panel might optionally include MUC 1 and/or c-myc, for colorectal cancer ras and/or APC , for prostate cancer PSA and/or BRCA 1 and/or BRCA2 or for ovarian
25 cancer BRCA1 and/or BRCA2 and/or CA125. There are other preferred embodiments in which p53 or c-erbB2 are not necessarily essential. For example, in the case of breast cancer suitable panels could be selected from the following:

30 p53 and MUC 1 with optional c-erbB2 and/or c-myc, and/or BRCA1 and/or BRCA2 and/or PSA;
p53 and c-myc with optional c-erbB2 and/or MUC1 and/or BRCA1 and/or BRCA2 and/or PSA;
35 p53 and BRCA1 with optional c-erbB2 and/or MUC 1 and/or c-myc and/or BRCA2 and/or PSA;
p53 and BRCA2 with optional c-erbB2 and/or MUC 1

and/or c-myc and/or BRCA1 and/or PSA;
c-erbB2 and MUC 1 with optional p53 and/or c-myc,
and/or BRCA1 and/or BRCA2 and/or PSA;
c-erbB2 and c-myc with optional p53 and/or MUC1 and/or
5 BRCA1 and/or BRCA2 and/or PSA;
c-erbB2 and BRCA1 with optional p53 and/or MUC 1
and/or c-myc and/or BRCA2 and/or PSA;
c-erbB2 and BRCA2 with optional p53 and/or MUC 1
and/or c-myc and/or BRCA1 and/or PSA;

10

In the case of colorectal cancer suitable panels
could be selected for example from the following:

p53 and ras with optional c-erbB2 and/or APC;

p53 and APC with optional c-erbB2 and/or Ras;

15

Ras and APC with optional p53 and/or c-erbB2

Such panels might also include CEA or CA19-9

In the case of prostate cancer suitable panels
could be selected for example from the following:

20

p53 and PSA with optional BRCA1 and/or BRCA2 and/or c-
erbB2;

c-erbB2 and PSA with optional p53 and/or BRCA1 and/or
BRCA2.

25

In the case of ovarian cancer suitable panels
could be selected for example from the following:

p53 and CA125 with optional c-erbB2 and/or BRCA1
and/or BRCA2;

c-erbB2 and CA125 with optional p53 and/or BRCA1

30

and/or BRCA2.

In a further embodiment, the immunoassay method
of the invention may be used in the selection of an
anti-cancer vaccine for use in a particular patient.

35

In this embodiment a sample of bodily fluid taken from
the patient is tested using a panel of two or more
immunoassay reagents, each corresponding to a

different tumour marker protein, in order to determine the relative strength of the patient's immune response to each of the different tumour marker proteins. The "strength of immune response" to a given tumour marker protein or proteins is indicated by the presence and/or the amount of cancer-associated autoantibodies specific to that tumour marker protein detected using the immunoassay; where autoantibodies are quantified, the greater the level of cancer-associated auto-antibodies, the stronger the immune response. The tumour marker protein or proteins identified as eliciting the strongest immune response or responses in the patient (i.e. the highest level of autoantibodies) is or are then selected to form the basis of an anti-cancer vaccine for use in the patient.

In a further embodiment, the invention provides a method of monitoring whether vaccination of a subject with an anti-cancer vaccine based on a particular tumour marker protein has been successful in eliciting a humoral immune response (i.e. antibodies against the said tumour marker protein). This method is based on the same immunoassay methodology used to measure cancer-associated anti-tumour autoantibodies (i.e. use of an immunoassay reagent based on tumour marker protein purified from a body cavity fluid or an excretion taken from a cancer patient), the only difference being what is measured in the assay is an antibody response rather than an autoantibody response.

In this embodiment a sample of bodily fluid taken from a patient previously treated with the anti-cancer vaccine (e.g. an immunogenic preparation comprising the relevant tumour marker protein, or an antigenic fragment thereof or a vaccine comprising a nucleic

acid encoding said relevant tumour marker protein) is contacted with an immunoassay reagent and complexes formed by specific binding of the immunoassay reagent to cancer-associated antibodies present in the sample are detected. The immunoassay reagent again comprises a sample of the said tumour marker protein prepared from bodily fluid derived from a body cavity or space as defined herein from one or more cancer patients and/or tumour marker protein prepared from an excretion from one or more cancer patients.

In addition to clinical applications in the detection of cancer, etc., the method of the invention may be used in any application where it is desired to test for the presence of cancer-associated anti-tumour autoantibodies. The method or tumour marker protein preparation of the invention may have applications in the laboratory as a research tool.

For example, it is possible for tumour marker proteins to have utility as therapeutic agents. The availability of large quantities of protein as provided by the bodily fluids defined herein allows pre-clinical and clinical testing, either *in vitro* or *in vivo* in humans or non-human animals, to determine efficacy of particular tumour marker proteins as therapeutic agents. Such a method would be applicable to each or all of the various tumour marker proteins described herein.

Another utility for tumour marker preparations of the invention is as a calibration material to be used in conjunction with the development of diagnostic tests for the presence of cancer or risk of cancer, which tests are based upon determination of the presence and/or level of any particular tumour marker protein in a clinical sample from a patient. The

tumour marker protein preparations of the invention can be used to construct calibration curves for such tests. In particular this aspect of the invention includes:

5 A method of calibrating an assay for measurement or detection of a given tumour marker protein in a clinical sample which method comprises the steps of:

 a) preparing at least two samples of a tumour
10 marker protein prepared according to the method of the invention, each of which comprises said given tumour marker protein and each of which has a different tumour marker protein concentration to each of the other said samples:

 b) carrying out a quantitative measurement of the
15 concentration of said tumour marker protein in each of said samples using:

 i) a spectrometric or spectrophotometric
 method and/or,
 ii) an antibody reagent to said tumour
20 marker protein, and

 c) constructing a standard curve for tumour
marker protein concentration based on the measurements obtained in step (b).

25 Such standard curves may be constructed for any or all of the specific tumour marker proteins described herein.

30 The invention will be further understood with reference to the following experimental Examples, together with the accompanying Figures in which:

 Figure 1 shows a post-Ig disruption gel
filtration chromatogram of a preparation of MUC16
(CA125) from ascites;

35 Figure 2 shows a silver stained gel of c-myc purification from ascitic fluid, post immunoaffinity chromatography;

Figure 3 shows an immunoprobed blot, c-myc purification from ascitic fluid, post immunoaffinity chromatography;

5 Figure 4 shows a comparison of patient serum (patients with no evidence of breast cancer themselves but with a family history of breast cancer and those with primary breast cancer) auto-antibody reactivity against MUC1 isolated from various body fluids: urine (from "normal" individuals), pleural effusion from a cancer patient and serum from advanced breast cancer patients (ABC serum);

10 Figure 5 shows autoantibody reactivity in serum from normal individuals against MUC1 from various body fluids: urinary MUC1 (normal), pleural effusion from a cancer patient and from advanced breast cancer patients (ABC serum);

15 Figure 6 shows the autoantibody reactivity in serum samples from pre-operative patients with ovarian masses against normal MUC16 (CA125) and against tumour-associated MUC16 from ascites;

20 Figure 7 shows the cancer-associated MUC1 concentration in sera, pleural effusion and ascitic fluid;

25 Figure 8 shows the cancer-associated MUC1 concentration in serum, wound drainage fluid and in seroma;

30 Figure 9 shows the reactivity of purified autoantibodies from seroma of patient M with cancer against purified urinary MUC1 from patient M taken two years prior to cancer diagnosis, MUC1 derived from the seroma of patient M, after diagnosis with cancer and bovine serum albumen conjugated to MUC1 protein core peptide;

35 Figure 10 shows serum autoantibody reactivity against MUC1 purified from pooled ascites fluid and against MUC1 purified from individual ascites samples from cancer patients;

Figure 11 shows serum autoantibody reactivity against MUC1 purified from pooled pleural effusions and against MUC1 purified from individuals pleural effusion samples from cancer patients; and

5 Figure 12 shows a calibration curve prepared from MUC1 from a pleural effusion.

Example 1-General protocol for purification of MUC1 antigen

- 10 Monoclonal anti-MUC1 antibody B55 (also known as NCRC 11, Xoma Corporation) is conjugated to CNBr-sepharose beads. Other anti-MUC1 monoclonal antibodies may be substituted for B55.
- 15 Tumour-induced body fluids (e.g. pleural effusion, ascites, seroma or wound drainage fluid) are diluted 1/10 with phosphate buffered saline (PBS) and filtered to 0.45 μ m.
- 20 Diluted body fluids are incubated with the anti-MUC1 sepharose beads (25 ml diluted fluid to 1 ml packed volume of beads) overnight at 4 °C with rolling ("batch" method) or re-circulated overnight through a packed column containing anti-MUC1 sepharose beads
- 25 ("column" method).

"Batch" method:-

30 Beads are packed by centrifugation and the supernatant removed;

Beads re-suspended in 5-10 ml PBS and rolled for 10 mins then packed by centrifugation and the supernatant removed; repeat 5 times (or until $A_{280\text{ nm}} \sim 0$);

35 Beads re-suspended in 5 ml 100 mM DEA pH11, and rolled at room temperature for 10 mins;

Beads packed by centrifugation and the supernatant removed, pH adjusted to 7 by the addition of pH 7 Tris buffer, dialysed against PBS for 24 hours minimum (100 DEA fraction);

5

Beads re-suspended in 5 ml PBS and rolled for 10 mins then packed by centrifugation and the supernatant removed, pH adjusted to 7 by the addition of pH 7 Tris buffer, dialysed against PBS for 24 hours minimum (post-DEA fraction);

10

MUC1 content of each fraction confirmed by ELISA using, for instance, the monoclonal anti-MUC1 antibody C595 (available from Cancer Research Campaign Laboratories, UK) (see example 5 for details) or B55, prior to pooling of the two fractions and storage at -20°C.

15

"Column" method:-

20

Column washed with 5 column volumes of PBS, or until eluate reads ~0 at $A_{280\text{ nm}}$;

25

1 column volume of 100 mM DEA pH11 applied, followed by 5 column volumes of PBS;

Eluate fractions (2ml) collected from the time of DEA application through the application of PBS;

30

Fractions dialysed overnight against PBS;

35

Fractions assayed for MUC1 content by ELISA using, for instance, the monoclonal anti-MUC1 antibody C595 or B55, prior to pooling MUC1 positive fractions and storage at -20°C.

In order to remove contaminating immunoglobulins, MUC1

pooled fractions are incubated with dithiothreitol (DTT) to 50 mM for 30 mins, then iodoacetamide (to 75 mM) before being subjected to gel filtration on an S300 column.

5

Resulting fractions (5ml) are assayed for MUC1 and human immunoglobulin (Ig) content by ELISA.

10 MUC1 containing fractions (uncontaminated with human Ig) are pooled and stored at -20°C.

Example 2a-General protocol for purification of MUC16 antigen (previously known as CA125)

15 One volume (e.g. 50 ml) of saturated ammonium sulphate was added to one volume (e.g. 50 ml) of tumour-induced body fluid (e.g. pleural effusion, ascites, seroma or wound drainage fluid) and incubated overnight at 4°C.

20 The resultant precipitate is collected by centrifugation (3500 rpm for 30 min in a standard benchtop centrifuge) and resuspended in ½ volume PBS.

25 This resuspension is subjected to gel filtration chromatography through an S300 column (2.5 x 100 cm) using PBS as the eluting buffer.

30 Fractions (5 or 10 ml) are collected and assayed by ELISA for MUC16, using for instance anti-CA125 from ICN or the anti-MUC16 antibody VK8 (Memorial Sloane Kettering, New York), prior to pooling MUC16 positive fractions and storage at -20°C.

35 In order to remove contaminating immunoglobulins, MUC16 pools are incubated with NaSCN (to 1.5M) for 10 mins, DTT (to 50mM) for 30 mins, then iodoacetamide (to 75mM) for 30 mins before being subjected to gel

filtration on, for instance, an S300 or a Superdex™ 75 column.

5 Resulting fractions (5ml) are assayed for MUC16 and human immunoglobulin (Ig) content by ELISA.

MUC16 containing fractions (uncontaminated with human Ig) are pooled and stored at -20°C.

10 Example 2b-Post Ig disruption gel filtration chromatography

For a sample prepared in the manner described above, fractions from a post-Ig disruption gel filtration
15 were assayed for MUC16 using anti-MUC16 antibody VK8 and for human Ig using an anti-human Ig. The results are shown in Figure 1. As is clearly demonstrated, two substantially immunoglobulin free MUC16 peaks are eluted.

20

Example 3-purification of c-myc antigen

Methodology as per purification of MUC1 (Example 1), except that:

25

Monoclonal anti-c-myc antibody 9E10 (ATCC) is used (or equivalent anti-c-myc antibody).

Gel filtration is performed on a Superdex™ 75 column.

30

Electrophoresis and Western blotting

Purity of MUC1, MUC16 and c-myc fractions are assessed by denaturing polyacrylamide gel
35 electrophoresis and Western blotting, performed according to standard protocols using BioRad™ Mini Protean III™ system and BioRad™ DryBlot™ system.

Protein patterns were revealed on gels for c-myc by silver staining (Figure 2). Western blots of c-myc were immuno-probed using monoclonal antibodies 9E10 (Figure 3). In each case, c-myc as well as
5 immunoglobulin heavy and light chains are identified.

Example 4-standard auto-antibody assay

10 Tumour antigen (e.g. MUC1, MUC16 or c-myc prepared according to Examples 1-3) diluted appropriately in PBS is plated out at 50 µl per well in a standard 96 well microtiter plate and left to air dry overnight;

15 Plate washed once with PBS/Tween™ to remove residual salt crystals;

Plate blocked for 60mins with 0.1% casein or 1% BSA in PBS;

20 Plate washed x3 with PBS/Tween™;

Serum (diluted 1/100 in PBS/0.1% casein) plated out in triplicate (50 µl per well), also monoclonal antibody controls;

25 Incubate for 60 mins at room temperature with shaking;

Wash plate x4 with PBS/Tween™;

30 Add horseradish peroxidase (HRP)-conjugated anti-Ig antibody (Dako) to each well (50 µl per well) at 1/8000 dilution for anti-human and 1/1000 for anti-mouse;

35 Incubate for 60 mins at room temperature with shaking;

Wash plate x4 with PBS/Tween™;

Add 50µl TMB (tetramethylbenzadine) per well and read kinetically over a 10 min period at $A_{650 \text{ nm}}$.

Experimental Data

5 Using the method as described in Example 4, cancer-associated autoantibodies to MUC1 and MUC16 were measured in a variety of sera using MUC1 and MUC16 isolated from the various sources as described herein. Results generated are shown in Figures 4 to 6.

10

Figure 4 shows a comparison of patient serum auto-antibody reactivity against MUC1 isolated from various body fluids: urine (from "normal" individuals), pleural effusion from a cancer patient and serum from advanced breast cancer patients (ABC serum). The patient serum tested was from either individuals with no evidence of breast cancer themselves but with a family history of breast cancer (i.e. one or more relatives who had breast cancer at a young age) or individuals with primary breast cancer.

20

Standard auto-antibody ELISAs were performed as described above, utilising MUC1 isolated from urine (normal), pleural effusion or ABC serum as antigen. Data was normalised to an internal control reaction using the DF3 anti-MUC1 monoclonal antibody (as opposed to a serum sample) against each of the MUC1 antigens.

25

30 As can be seen from the Figure, MUC1 derived from normal urine (nMUC1) was consistently lower in its reactivity than MUC1 derived from either pleural effusion (PE) or ABC serum. Furthermore, MUC1 derived from PE was of similar reactivity to cancer-associated MUC1 autoantibodies as MUC1 isolated from the serum of patients with ABC and therefore of equal diagnostic value.

35

Figure 5 shows the results of an identical exercise to Figure 4 except that all serum samples tested were for normal individuals (no breast cancer or family history of breast cancer). As can be seen, there is no significant difference in the reactivity of the serum to the three different antigens.

Figure 6 shows reactivity of MUC16 cancer-associated autoantibodies from serum of patients with ovarian masses (pre-operative) against MUC16 (CA125) isolated from the serum of normal individuals and from ascites fluid in a patient with breast cancer. Antigens were prepared as in Example 2 and autoantibodies detected using ELISA assay as described in Example 4.

As can be seen, greatly enhanced reactivity of the cancer-associated MUC16 autoantibodies is seen with the MUC16 antigen from ascites fluid as compared to the 'normal' MUC16. This experimental result therefore confirms the usefulness of ascites fluid as an antigen source for detection of cancer-associated autoantibodies.

Example 5-Measurement of cancer-associated MUC1 levels in ascites fluid, pleural effusion, seroma and wound drainage fluid

MUC1 levels found in the serum of a patient with cancer were compared with the levels found in ascites fluid, pleural effusion, wound drainage fluid or seroma, in each case in the same patient from whom the serum sample was taken. MUC1 in the samples was quantified according to the following protocol:

Capture MUC1 ELISA Protocol

Aliquot 50 μ l well⁻¹ antibody solution into triplicate

wells of a microtitre plate (usually $1\mu\text{g ml}^{-1}$ C595(IgG) and appropriate negative control) and incubate at RT with shaking for 1hr for the protein to adsorb to the plate.

5

Wash the plate x4 with PBS/Tween using $250\mu\text{l well}^{-1}$.

Block the plate using 1% BSA $100\mu\text{l well}^{-1}$ and incubate at RT with shaking for 1hr.

10

Wash the plate x4 with PBS/Tween using $250\mu\text{l well}^{-1}$.

Apply $50\mu\text{l}$ per well of fluid being tested, diluted 1/10 in PBS and incubate at RT with shaking for 1hr.

15

Wash the plate x4 with PBS/Tween using $250\mu\text{l well}^{-1}$.

Add $50\mu\text{l well}^{-1}$ biotinylated C595 ($1\mu\text{g/ml}$) and incubate at RT with shaking for 1hr.

20

Wash the plate x4 with PBS/Tween using $250\mu\text{l well}^{-1}$.

Add $50\mu\text{l well}^{-1}$ extra-avidin peroxidase at 1/1000 dilution and incubate at RT with shaking for 1hr.

25

Wash the plate x4 with PBS/Tween using $250\mu\text{l well}^{-1}$.

Add $50\mu\text{l well}^{-1}$ TMB substrate and read kinetically at 650_{nm} for 10 minutes.

30

The results are shown in Figures 7 and 8.

As will be readily apparent from the data serum levels of the cancer-associated MUC1 antigen are significantly lower than the level found in either ascites fluid, pleural effusion, seroma or wound drainage fluid. Accordingly, there are substantial

35

benefits to be gained in terms of yield in recovering tumour marker antigen from those body cavity fluids.

Example 6-Reactivity of human anti-MUC1 antibodies
5 purified against cancer-associated MUC1 from seroma

Human antibodies from seroma from patient M were purified by immunoaffinity chromatography against MUC1 derived from seroma fluid from the same cancer patient
10 M. Purified antibodies were then tested against BSA conjugated protein core peptide to MUC1 and MUC1 derived from:- patient M's urine taken two years prior to cancer diagnosis; patient M's seroma taken after cancer diagnosis. The antibody purification from
15 seroma was carried out according to the following protocol:

Human anti-MUC1 antibody purification

20 Purification of human anti-MUC1 auto-antibodies was by affinity chromatography.

Seroma fluid, diluted 10 fold in PBS pH 7.6, was applied at 0.5ml/min by overnight re-circulation at
25 4°C, to an affinity matrix in column format, consisting of CNBr sepharose (Pharmacia) coupled (following the manufacturers instructions) to Pt-MUC1.

After seroma fluid application, the column was washed
30 with 15ml of PBS (ensuring return of $A_{280\text{nm}}$ reading to zero) prior to elution of antibody using 10ml of 3M NaSCN, at 1ml/min.

Fractions of 1ml were collected throughout, desalted
35 by dialysis against PBS and tested by ELISA for the presence of antibody.

Positive fractions were pooled, purity of antibody verified (by PAGE) and antibody concentration determined.

- 5 Assay of the purified antibodies against the three MUC1 antigens identified above was carried out according to the following protocol:

MUC1 ELISA Protocol

10

Aliquot 50 μ l well⁻¹ of the MUC1 antigen solution into triplicate wells of a microtitre plate and dry down at RT overnight.

15

Wash the plate x2 with PBS/Tween using 250 μ l well⁻¹.

Block the plate with 1% BSA using 100 μ l well⁻¹ and incubate at RT with shaking for 1hr.

20

Wash the plate x2 with PBS/Tween using 250 μ l well⁻¹.

Add 50 μ l well⁻¹ purified antibody solution at 1 μ g/ml and incubate at RT with shaking for 1hr.

25

Wash the plate x4 with PBS/Tween using 250 μ l well⁻¹.

Add 50 μ l well⁻¹ α -human Ig HRP (DAKO), freshly diluted as per manufacturers instructions, and incubate at RT with shaking for 1hr.

30

Wash the plate x4 with PBS/Tween using 250 μ l well⁻¹.

Add 50 μ l well⁻¹ TMB substrate and read kinetically at 650_{nm} for 10 minutes.

35

The results are shown in Figure 9.

Reactivity of the antibodies against MUC1 peptide was negligible. Reactivity of antibodies against normal MUC1 was considerably lower than that seen towards patient M's seroma derived MUC1. It can be inferred from this result that normal MUC1 molecule is substantially different with regard to its immune recognition, to that found in seroma fluid from an individual with cancer.

Example 7-Serum reactivity against MUC1 purified from pooled ascitic fluid and pleural effusions

MUC1 was purified from pooled ascitic fluid and from pooled pleural effusion from patients with advanced breast cancer using the protocol described in Example 1 and its reactivity against serum from patients with primary breast cancer measured as described in Example 4. The antigen from the pooled fluids was compared in each case with antigen isolated from 3 individual samples of ascitic fluid or pleural effusion respectively from patients with ABC. The results are shown in Figures 10 and 11.

In the case of both ascitic fluid and pleural effusion the reactivity of the MUC1 from pooled fluid is as good as that isolated from individual samples. Furthermore, while there is great scope for variability of reactivity using samples from individuals, pooled samples provide greater consistency of product so that one would not expect the reactivity to significantly vary between batches from pooled samples.

Example 8-Calibration Curve using MUC1

Serial dilutions of MUC1 which had been isolated from pleural effusion were prepared. Their MUC1

concentrations were measured by the method as shown in
example 4 except that no human sera were used.
Detection was by mouse B55 antibody followed by Dako
anti-mouse HRP using an end-point rather than a
5 kinetic reading.

The results are shown in Figure 12 and confirm the
utility of the tumour marker proteins prepared in
accordance with the invention as a calibration
10 material.

Claims

1. A method of detecting cancer-associated anti-tumour autoantibodies, which method is an immunoassay comprising contacting a sample to be tested for the presence of such autoantibodies with an immunoassay reagent and detecting the presence of complexes formed by specific binding of the immunoassay reagent to any cancer-associated anti-tumour autoantibodies present in the sample, wherein the immunoassay reagent comprises tumour marker protein prepared from bodily fluid, derived from a body cavity or space in which a tumour is or was present or with which a tumour is or was associated, of one or more cancer patients and/or tumour marker protein prepared from an excretion of one or more cancer patients wherein said tumour marker protein exhibits selective reactivity with cancer-associated anti-tumour autoantibodies.

2. A method according to claim 1 which comprises performing an immunoassay to detect and/or quantitatively measure the presence of two or more types of autoantibodies, each immunologically specific to different tumour marker proteins or to two or more epitopes of the same tumour marker protein, wherein the immunoassay is carried out using a panel of two or more immunoassay reagents, at least one of which reagents comprises tumour marker protein prepared from bodily fluid derived from a body cavity or space from one or more cancer patients and/or tumour marker protein prepared from an excretion from one or more cancer patients.

3. Use of the method of claim 1 or claim 2 for the detection or diagnosis of cancer in a patient, wherein the sample to be tested using the immunoassay

is a sample of bodily fluid from taken from the patient, and wherein the presence of an elevated level of autoantibodies, as compared to normal control individuals, is taken as an indication that the individual has or is developing cancer.

4. Use of the method of claim 1 or claim 2 in monitoring the progress of cancer or other neoplastic disease in a patient, wherein the sample to be tested using the immunoassay is a sample of bodily fluid taken from the patient, and wherein the presence of an elevated level of autoantibodies, as compared to a normal control, is taken as an indication of the presence of cancer in the patient.

5. Use of the method of claim 1 or claim 2 in detecting early neoplastic or early carcinogenic change in an asymptomatic subject, wherein the sample to be tested using the immunoassay is a sample of bodily fluid taken from the subject, and wherein the presence of an elevated level of autoantibodies, as compared to normal control individuals, is taken as an indication of early neoplastic or early carcinogenic change in the subject.

6. Use of the method of claim 1 or claim 2 in screening a population of asymptomatic human subjects to identify those subjects who are at increased risk of developing cancer, wherein the samples to be tested using the immunoassay are samples of bodily fluid taken from the subjects, and wherein subjects having an elevated level of autoantibodies, as compared to normal control individuals, are identified as being at risk of developing cancer.

7. Use of the method of claim 1 or claim 2 in monitoring the response of a cancer patient to anti-

cancer treatment, wherein the sample to be tested using the immunoassay is a sample of bodily fluid taken from the patient, and wherein the presence of a decreased level of autoantibodies after treatment is taken as an indication that the patient has responded positively to the treatment.

8. Use of the method of claim 1 or claim 2 in the detection of recurrent disease in a patient previously diagnosed as having cancer, which patient has undergone anti-cancer treatment to reduce the amount of cancer present, wherein the sample to be tested using the immunoassay is a sample of bodily fluid taken from the patient, and wherein the presence of an increased level of autoantibodies in the patient, as compared to a normal control, is taken as an indication that disease has recurred.

9. Use of the method of claim 2 in the selection of an anti-cancer vaccine for use in a particular patient, wherein the immunoassay is carried out using a panel of two or more immunoassay reagents each corresponding to a different tumour marker protein in order to determine the relative strength of the patient's immune response to each of the different tumour marker proteins, wherein the tumour marker protein or proteins identified as eliciting the strongest immune response or responses in the patient is or are selected to form the basis of an anti-cancer vaccine for use in said patient.

10. A method of determining whether a vaccination procedure comprising challenging a patient with an immunogenic preparation comprising a tumour marker protein or an antigenic fragment thereof or with a nucleic acid sequence expressing said tumour marker protein, has been successful in eliciting

cancer-associated antibodies to the tumour marker protein in the patient, which method is an immunoassay comprising contacting a sample of bodily fluid from the patient with an immunoassay reagent and detecting
5 the presence of complexes formed by specific binding of the immunoassay reagent to any cancer-associated antibodies present in the sample, wherein the immunoassay reagent comprises a sample of the said tumour marker protein prepared from bodily fluid
10 derived from a body cavity or space in which a tumour is or was present or with which a tumour is or was associated from one or more cancer patients and/or tumour marker protein prepared from an excretion from one or more cancer patients, wherein said tumour
15 marker protein exhibits selective reactivity with cancer-associated anti-tumour antibodies.

11. A method according to any one of claims 1, 2 or 10 wherein the bodily fluid derived from a body
20 cavity or space is ascites fluid, pleural effusion, seroma, hydrocoele or wound drainage fluid.

12. The use according to anyone of claims 3 to 9 wherein the bodily fluid derived from a body cavity or
25 space is ascites fluid, pleural effusion, seroma, hydrocoele or wound drainage fluid.

13. A method according to any one of claims 1, 2 or 10 wherein the excretion is urine, faeces or
30 seminal fluid.

14. The use according to any one of claims 3 to 9 wherein the excretion is urine, faeces or seminal
fluid.

35 15. A method according to claim 11 or 13 wherein the tumour marker protein is selected from MUC1, MUC16

or c-myc.

16. The use according to claim 12 or 14 wherein
the tumour marker protein is selected from MUC1, MUC16
5 or c-myc.

17. A method according to any one of claims 1,
2, 10, 11 or 13 wherein the tumour marker protein is
selected from c-erbB2, p53, ras, BRCA1, BRCA2, APC,
10 PSA, CEA, and CA19.9.

18. The use according to any one of claims 3 to
9, 12 or 14, wherein the tumour marker protein is
selected from c-erbB2, p53, ras, BRCA1, BRCA2, APC,
15 PSA, CEA and CA19.9.

19. Use of tumour marker protein prepared from
bodily fluid derived from a body cavity or space in
which a tumour is or was present or with which a
20 tumour is or was associated, of one or more cancer
patients and/or tumour marker protein derived from an
excretion of one or more cancer patients in the
manufacture of an immunoassay reagent exhibiting
selective reactivity with cancer-associated anti-
25 tumour autoantibodies.

20. A method of preparing a tumour marker
protein which method comprises isolating said tumour
marker protein from bodily fluid wherein said fluid
30 is:

- (i) collected from a body cavity or space
in which a tumour is or was present or
with which a tumour is or was
associated, and
- 35 (ii) said fluid represents the pooled fluid
samples from two or more cancer
patients.

21. A method as claimed in claim 20 wherein said fluid is acites, pleural effusion, seroma, hydrocoele or wound drainage fluid or a mixture thereof.

5 22. A method as claimed in claim 21 wherein said tumour marker protein is MUC1.

10 23. A method as claimed in claim 20 or 21 wherein said tumour marker protein is c-erbB2, p53, ras, BRCA1, BRCA2, APC, PSA, CEA, CA19.9, MUC16 or c-myc.

15 24. A method as claimed in any one of claims 20 to 23 wherein said tumour marker is purified from said fluid by affinity chromatography.

20 25. A method as claimed in any one of claims 20 to 24 which includes a step of removing contaminating immunoglobulin from said tumour marker protein.

 26. A method as claimed in any one of claims 20 to 25 wherein said tumour marker protein is immobilized to solid support.

25 27. A method of preparing a tumour marker protein which method comprises isolating said tumour marker protein from a bodily fluid collected from a body cavity or space in which a tumour is or was present or with which a tumour is or was associated
30 wherein said bodily fluid is wound drainage fluid, seroma, hydrocoele or a mixture thereof.

35 28. A method of preparing a tumour marker protein which method comprises isolating said tumour marker protein from an excretion from one or more cancer patients, said excretion or any component thereof, having been in contact with a tumour or

tumour cells.

29. A method as claimed in claim 28 wherein said excretion is urine, faeces or seminal fluid.

5

30. A method as claimed in claim 28 or 29 wherein the relevant component of said excretion is bile.

10

31. A method as claimed in any one of claims 27 to 30 including any one of the specific features defined in claims 22 to 26.

15

32. A preparation of a tumour marker protein prepared by the method of any one of claims 20 to 31 and which is substantially immunoglobulin free.

20

33. A kit or reagent suitable for carrying out an immunoassay which comprises a preparation of a tumour marker protein as claimed in claim 31 immobilized to a solid support.

25

34. A kit or reagent as claimed in claim 33 wherein said solid support is the surface of a well of a multiwell plate or is a bead.

30

35. A kit or reagent as claimed in claim 32 or 33 wherein said immobilized tumour marker protein is absorbed or covalently attached to said solid support.

35

36. Use of a preparation as claimed in claim 32 in the evaluation in an *in vitro* test for the therapeutic efficacy or safety of said tumour marker protein.

37. Use of a preparation as claimed in claim 32 in manufacture of a composition for the evaluation in

an *in vivo* test of the therapeutic efficacy or safety of said tumour marker protein.

5 38. A method of calibrating an assay for measurement or detection of a given tumour marker protein in a clinical sample which method comprises the steps of:

10 a) preparing at least two samples of a preparation of claim 32, each of which comprises said given tumour marker protein and each of which has a different tumour marker protein concentration to each of the other said samples:

15 b) carrying out a quantitative measurement of the concentration of said tumour marker protein in each of said samples using

 (i) a spectrophotometric method and/or,

 (ii) an antibody reagent to said tumour marker protein, and

20 c) constructing a standard curve for a tumour marker protein concentration based on the measurements obtained in step (b).

25



INVENTOR IN PEOPLE

Application No: GB 0226622.9**Examiner:**

Dr Rowena Dinham

Claims searched: 20-27; and 31-38 (in part)**Date of search:**

29 July 2003

Patents Act 1977 : Further Search Report under Section 17**Documents considered to be relevant:**

Category	Relevant to claims	Identity of document and passage or figure of particular relevance
X	20-26 & 31-38	<i>Mol Immunol</i> ; Vol 18, pp 647-653 (1981). Hill <i>et al</i> "Nature of carinoembryonic antigen purified from malignant ascitic fluid..." See entire document, especially Materials and Methods "Sephadex gel chromatography" and Figure 1
X	20-26 & 31-38	<i>Clin Cancer Res</i> ; Vol 7, pp 781s-787s (2001). Beatty <i>et al</i> . "Biochemical characterization of the soluble form of tumor antigen MUC1..." See entire document, especially Materials and Methods 'ASC-MUC1 purification'
X	20-26 & 31-38	WO 92/11281 A1 (NEW ENGLAND DEACONESS HOSPITAL CORP) See entire document, especially Examples 1 and 2
X	20-26 & 31-38	<i>Br J Cancer</i> ; Vol 36, pp 1046-1052 (1978). Wolf. "A tumour associated antigen from the pleural effusion..." See entire document, especially Figure 2
X	20-26 & 31-38	<i>Biochem Biophys Res Commun</i> ; Vol 171, pp 633-640 (1990). Toth <i>et al</i> . "A carcinoembryonic antigen (CEA) binding protein from ascites..." See entire document, especially Materials and Methods "CEA purification and analysis"
X	20-26 & 31-38	EP 1189063 A1 (HEALTH RESEARCH INC) See entire document, especially page 5, para [0027]

Categories:

X	Document indicating lack of novelty or inventive step	A	Document indicating technological background and/or state of the art
Y	Document indicating lack of inventive step if combined with one or more other documents of same category.	P	Document published on or after the declared priority date but before the filing date of this invention.
&	Member of the same patent family	E	Patent document published on or after, but with priority date earlier than, the filing date of this application.

Field of Search:Search of GB, EP, WO & US patent documents classified in the following areas of the UKC^v:

--

Worldwide search of patent documents classified in the following areas of the IPC⁷:

--



INVESTOR IN PEOPLE

Application No: GB 0226622.9

Examiner: Dr Rowena Dinham

Claims searched: 20-27; and 31-38 (in part)

Date of search: 29 July 2003

G01N

The following online and other databases have been used in the preparation of this search report :

WPI, EPODOC, JAPIO, MEDLINE, EMBASE, BIOSIS, SCISEARCH, CAPLUS



INVESTOR IN PEOPLE

Application No: GB 0226622.9 **Examiner:** Dr Rowena Dinham
Claims searched: 28-30; and 31-38 (in part) **Date of search:** 29 July 2003

Patents Act 1977 : Further Search Report under Section 17

Documents considered to be relevant:

Category	Relevant to claims	Identity of document and passage or figure of particular relevance
X	28, 29 & 31-38	<i>Biochim Biophys Acta</i> ; Vol 1336, pp 425-433 (1997). Shibata <i>et al.</i> "Purification and characterization of prostate specific antigen from human urine." See entire document, especially Materials and methods 2.2 "Purification of prostate specific antigen"
X	28, 29 & 31-38	<i>Clin Chem</i> ; Vol 34, pp 2528-2532 (1988). Rusciano <i>et al.</i> "Concomitant purification of prostatic carcinoma tumour markers..." See entire document, especially Materials and Methods
X	28, 29 & 31-38	<i>Br J Cancer</i> ; Vol 29, pp 413-424 (1974). Nery <i>et al.</i> "Isolation and partial characterisation of macromolecular urinary aggregates..." See entire document, especially Materials and Methods
X	28, 29 & 31-38	EP 1189063 A1 (HEALTH RESEARCH INC) See entire document, especially page 5, para [0027] and page 7, para [0047]-[0048]
X	28, 30 & 31-38	<i>Sao Paulo Med J</i> ; Vol 119, pp 110-113 (2001). Aparecida <i>et al.</i> "Value of CEA level determination in gallbladder bile..." See entire document, especially Results
X	28, 30 & 31-38	<i>J Biol Chem</i> ; Vol 269, pp 14430-14437 (1994). Baechström <i>et al.</i> "Purification and characterization of sialyl-Le ^a -carrying mucins of human bile..." See entire document, especially Results
X	28, 29 & 31-38	<i>Eur J Cancer</i> ; Vol 13, pp 433-436 (1987). Stubbs <i>et al.</i> "Faecal carcinoembryonic antigen in patients with large bowel cancer". See entire document, especially Results
X	28, 29 & 31-38	WO 00/19208 A1 (UNIVERSITY OF ULSTER) See entire document, especially Experiment 1
X	28, 29 & 31-38	<i>The Prostate</i> ; Vol 43, pp 150-157 (2000). Sokoloff <i>et al.</i> "A dual-monoclonal sandwich assay for prostate-specific

Categories:



INVESTOR IN PEOPLE

Application No: GB 0226622.9

Examiner: Dr Rowena Dinham

Claims searched: 28-30; and 31-38 (in part)

Date of search: 29 July 2003

X	Document indicating lack of novelty or inventive step	A	Document indicating technological background and/or state of the art
Y	Document indicating lack of inventive step if combined with one or more other documents of same category	P	Document published on or after the declared priority date but before the filing date of this invention.
&	Member of the same patent family	E	Patent document published on or after, but with priority date earlier than, the filing date of this application

Field of Search:

Search of GB, EP, WO & US patent documents classified in the following areas of the UKC^v:

--

Worldwide search of patent documents classified in the following areas of the IPC⁷:

G01N

The following online and other databases have been used in the preparation of this search report :

WPI, EPODOC, JAPIO, MEDLINE, EMBASE, BIOSIS, SCISEARCH, CAPLUS
--



INVENTOR IN PEOPLE

Application No: GB 0226622.9
Claims searched: 1-19

Examiner: Dr Rowena Dinham
Date of search: 13 May 2003

Patents Act 1977 : Search Report under Section 17

Documents considered to be relevant:

Category	Relevant to claims	Identity of document and passage or figure of particular relevance
Y	1 at least	WO 99/58978 A2 (UNI OF NOTTINGHAM) See especially page 5 line 2-22 and Example 5(C)
Y	1 at least	<i>Clin Cancer Res</i> ; Vol 7, pp 781s-787s (2001). Beatty <i>et al.</i> "Biochemical characterisation of the soluble form..." See especially Materials and Methods 'ACS-MUC1 purification'
Y	1 at least	<i>Lung</i> ; Vol 170, pp 65-74 (1992). Lawniczak <i>et al.</i> "The search for tumour-associated proteins in pleural effusions..." See especially Materials and Methods and Results
Y	1 at least	<i>Clin Chim Acta</i> ; Vol 300, pp 43-55 (2000). Kuralay <i>et al.</i> "Diagnostic usefulness of tumour marker levels..." See especially Materials and Methods and Results
A		WO 00/26668 A2 (UNI OF MICHIGAN) See especially page 6 line 7-20 and Examples
A		WO 01/11372 A1 (UNI OF MICHIGAN) See especially page 8 line 3-18 and Examples
A		<i>Intern Med</i> ; Vol 36, pp 724-727 (1997). Yamamoto <i>et al.</i> "L-Myc overexpression and detection of auto-antibodies..." See especially Materials and Methods 'Detection of anti-L-Myc antibodies' and 'Detection of L-myc antigen' and Results
A		<i>Int J Oncology</i> ; Vol 13, pp 605-610 (1998). Monternarh <i>et al.</i> "p53 autoantibodies in the sera, cyst and ascitic fluids..." See especially Materials and Methods 'ELISA procedure...' and Results
A		<i>Br J Cancer</i> ; Vol 80, pp 1987-1994 (1999). Schneider <i>et al.</i> "p53 protein, EGF receptor and anti-p53 antibodies..." See especially Patients and Methods 'Serum specimens' and Results
A		<i>Clin Cancer Res</i> ; Vol 4, pp 3025-3030 (1998). Lai <i>et al.</i> "Presence of serum anti-p53 antibodies..." See especially Materials and Methods 'p53 antigen' and Results

Categories:



INVESTOR IN PEOPLE

Application No: GB 0226622.9
Claims searched: 1-19

Examiner: Dr Rowena Dinham
Date of search: 13 May 2003

X	Document indicating lack of novelty or inventive step	A	Document indicating technological background and/or state of the art
Y	Document indicating lack of inventive step if combined with one or more other documents of same category.	P	Document published on or after the declared priority date but before the filing date of this invention.
&	Member of the same patent family	E	Patent document published on or after, but with priority date earlier than, the filing date of this application

Field of Search:

Search of GB, EP, WO & US patent documents classified in the following areas of the UKCV:

--

Worldwide search of patent documents classified in the following areas of the IPC⁷:

G01N

The following online and other databases have been used in the preparation of this search report:

WPI, EPODOC, JAPIO, MEDLINE, EMBASE, BIOSIS, SCISEARCH, CAPLUS
--